



# *New Concepts in Tree Planting*

*Bruce R. Fraedrich, Ph. D., Plant Pathologist*


**S**uccessful establishment of new plantings is dependent on correctly matching species to site and creating a soil environment that will allow new transplants to rapidly regenerate a root system. During transplanting, more than 95% of a root system is lost when the tree is dug by hand or with a tree spade. Soil surrounding the new plantings must have sufficient water, air space and nutrients to support rapid root regeneration to ensure plant survival. The following 10 steps will help ensure planting success.

**1. Assess planting site characteristics prior to species selection.** Soil analysis for pH, nutrient levels, organic matter content texture and drainage is essential to match species with the site. Attempting to alter soil conditions to support a species cultural preferences can be expensive and is seldom successful. The presence of utility lines, buildings, streets and other structures must be considered when selecting species that are suited to the site. A computer based tree selection program has been developed that assists in selecting species, which are compatible with site characteristics (*Southern Trees* 800-627-3819).

**2. Determine the available soil volume for the site** and select species and planting distances that allow the plant to reach maturity. Most premature decline of urban plantings is the result of water deficiency related to insufficient soil volume (root space). Recent research at Cornell University has provided methods to

determine soil volumes to meet the water demands of mature urban trees. In general, at least two cubic feet of soil is required for each square foot of crown projection (crown projection is the soil area beneath the crown). Calculating available soil volumes and selecting species whose mature size is compatible with the site is an integral component of the planning stage.

**3. Specify small caliper trees when possible.** Small caliper nursery stock regenerates a root system to pre-transplant levels more quickly than larger transplants. This translates into higher survival rates and lower maintenance costs during the establishment period. Crown growth rates of small transplants are much faster than larger stock following transplanting. After a ten-year period, two-inch transplants will be a similar



size to ten-inch stock. In general, 2 - 2 1/2" caliper or smaller stock is recommended. Larger sizes may be necessary in high use areas where vandalism is a factor.

**4. Prepare a large planting area.** The planting hole should be three-to-five times wider than the root ball especially in heavy clay or compacted soils. This will allow rapid regeneration of the root system following planting. The depth of the planting hole must be no greater than the root ball to prevent settling and root mortality. When preparing the planting site, amend the existing soil rather than use different backfill. In clay soils, using a well-drained loam backfill will result in water collecting in the planting hole following high rainfall, which can result in root mortality. Amend the soil with fertilizer, organic matter, lime or sulfur per soil analysis results. After planting cover the root ball and backfill with 2 - 4" of organic **mulch** such as wood chips to conserve soil moisture and moderate soil temperature. When planting in compacted or heavy clay soils, a large planting hole should be dug or the soil outside the hole should be tilled. After planting, mulch the entire area.

**5. Ensure the Root Collar of the transplant is flush with or slightly above the natural grade.** "Planting too deep" is a leading cause of death of new transplants. Specifications traditionally have required that the top of the root ball be flush with grade. In recent years, trees from some nurseries have soil on top of the root collar and against stem tissues. In these instances, the soil must be removed from the top of the ball until the root collar is exposed. Soil on top of the root collar will increase the incidence of stem disease and root disorders including root disease and girdling roots. When mulching after transplanting, make sure the mulch does not contact the trunk.


**6. Remove "foreign" materials from root balls.** Wire baskets must be removed from the upper 6-8 inches of the ball to prevent girdling of major support roots. Synthetic burlap, nylon twine and straps must be removed entirely from the root ball. Natural burlap can become water repellent and must be removed from the upper part of the ball to facilitate water infiltration.

**7. Do not wrap the trunk.** Recent research reveals that tree wraps provide little benefit in terms of stem protection from temperature extremes. Wraps may actually increase the incidence of certain pest infestations. Leave lower limbs to shade the trunk rather than wrapping.

**8. Stake or guy the plant only if necessary** to provide support or protect against strong winds and vandalism. Inspect guys periodically to ensure against girdling and bark injuries. Guys usually can be removed after one year.

**9. Prune only broken, dead and diseased limbs at planting time.** Research has shown that pruning the crown to "compensate for root loss" actually impedes root regeneration and slows establishment. Structural pruning should be delayed until the tree is established.

**10. Implement an intensive plant health care program following installation.** Young transplants stressed by root loss are sensitive to environmental stress and pest infestations. Periodically monitoring the trees for soil moisture conditions, decline symptoms, pest infestations, nutrient deficiencies and other plant health related conditions will facilitate early detection and correction of problems before mortality occurs.



*Mulching after transplanting will reduce water loss, moderate soil temperature and eliminate turf competition. These benefits should result in higher survival rate, faster growth, and a healthier tree.*

**Literature Cited**

1. Appleton, B.L. and S. French. 1992. *Current Attitudes Toward and Use of Tree Trunk Protective Wraps, Paints and Devices*. Journal of Arboriculture 18 (1): 15-19.
2. Gilman, E.F. et al. 1993. *Southern Trees: An Expert System for Selecting Trees*. Contact Robin Morgan, USFS, Atlanta, Georgia.
3. Harris, R.W., T. T. Leise and W.B. Davis. 1982. *Staking and Landscape Trees*. University of California Leaflet 2576. 13 pp.
4. Himelick, E.B. 1992. *Tree and Shrub Transplanting Manual* International Society of Arboriculture. 76 pp.
5. Lindsey, P. and N. Bassuk. 1991. *Specifying Soil Volumes to Meet the Water Demands of Mature Urban Street Trees and Trees in Containers*. Journal of Arboriculture 17 (6): 141-149.
6. McNeil, R. E., W.M. Fountain and M.L. Witt. 1982. *Tree Decline in the Landscape When the Trees were Dug B & B and plastic twine or wire baskets were used*. In Proc Urban Trees and Forests: Pest Management Problems, Needs and Prospects.
7. Shoup, S., R. Reavis and C.E. Whitcomb. 1981. *Effects of Pruning and Fertilizers on Establishment of Bareroot Deciduous Trees*. Journal of Arboriculture 7 (6): 155-157.
8. Smiley, E.T. 1991. *Root Collar Disorders*. Arbor Age 11(12): 40-41.
9. Smiley, E.T. 1991. *National Epidemic Reported: Improper Planting is Killing Trees* Arbor Age 11(12): 38.
10. Watson, G. 1985. *Tree Size Affects Root Regeneration and Top Growth After Transplanting*. Journal of Arboriculture 11: 37-40.